


AMENDMENTS TO THE CLAIMS









1. (Currently Amended) A cobalamin derivative of formula (I):



wherein:

- (i) R^b is a spacer-chelator group optionally carrying a metal atom;
- (ii) R^c, R^d, and R^e, and R^R are an antibiotic or antiproliferative therapeutic agent, each connected through a linker Z, or hydrogen; and R^R is an antibiotic or antiproliferative therapeutic agent connected through a linker Z, or hydrogen;
- (iii) with the proviso that at least one of the residues R^c, R^d, R^e and R^R are hydrogen;
- (iv) X is cyano, methyl, hydroxy, aquo or a 5'-deoxyadenosyl group; and

(v) the central cobalt (Co) atom is optionally in the form of a radioactive isotope; and wherein the spacer-chelator group consists of an aliphatic chain of 2 to 4 carbon atoms carrying a chelator selected from the chelators of formulae (II) to (IX):



wherein carboxyl groups in formulae (II) to (IX) may be present as esters; and said cobalamin derivative:

- (a) has no binding affinity or less than 20% binding affinity to transcobalamin II when compared to the binding affinity of non-modified cobalamin in a binding test, and
- (b) retains activity as a vitamin B12 substitute.

2. (Previously Presented) The cobalamin derivative according to claim 1 retaining more than 2% of the activity as a vitamin B12 substitute in a growth assay.

3. (Original) The cobalamin derivative according to claim 1

- (a) having less than 10% of binding affinity to transcobalamin II when compared to the binding affinity of non-modified cobalamin in a binding test, and
- (b) retaining more than 10% of the activity as a vitamin B12 substitute in a growth assay.

4. (Original) The cobalamin derivative according to claim 1

- (a) having less than 5% of binding affinity to transcobalamin II when compared to the binding affinity of non-modified cobalamin in a binding test, and
- (b) retaining more than 10% of the activity as a vitamin B12 substitute in a growth assay.

5. (Previously presented) The cobalamin derivative according to claim 1 carrying a therapeutic and/or diagnostic agent.

6. (Previously presented) The cobalamin derivative according to claim 1 carrying a radioactive metal.

7-10. (Cancelled)

11. (Previously presented) The cobalamin derivative according to claim 6 wherein the radioactive metal is ^{94m}Tc , ^{99m}Tc , ^{188}Re , ^{186}Re , ^{111}In , ^{90}Y , ^{64}Cu , ^{67}Cu or ^{177}Lu .

12. (Cancelled)

13. (Previously Presented) The cobalamin derivative according to claim 1 wherein X is cyano.

14. (Previously Presented) The cobalamin derivative according to claim 1, wherein the central cobalt atom is the radioisotope ^{57}Co or ^{60}Co .

15. (Previously Presented) The cobalamin derivative according to claim 1, wherein R^b is a spacer-chelator group optionally carrying a metal atom, the spacer is an aliphatic chain of 2 to 4 carbon atoms, and the chelator is of formula (II), wherein the group COOH is optionally in the form of an ester;
 R^c , R^d , R^e and R^R are hydrogen; and
X is cyano.

16. (Previously Presented) The cobalamin derivative according to claim 15, wherein R^b is a spacer-chelator group optionally carrying a metal atom, the spacer is an aliphatic chain of 4 carbon atoms, and the chelator is of formula (II), wherein the group COOH is in the form of the ethyl ester;
 R^c , R^d , R^e and R^R are hydrogen; and
X is cyano.

17. (Previously Presented) The cobalamin derivative according to claim 1, wherein R^d is a spacer-chelator group optionally carrying a metal atom, the spacer is an aliphatic chain of 3 carbon atoms, and the chelator is of formula (II), wherein the group COOH is optionally in the form of an ester;
 R^b , R^c , R^e and R^R are hydrogen; and
X is cyano.

18. (Previously Presented) The cobalamin derivative according to claim 1, wherein R^b is a spacer-chelator group optionally carrying a metal atom, the spacer is an aliphatic chain of 2 carbon atoms, and the chelator is of formula (III);
 R^c , R^d , R^e and R^R are hydrogen; and
X is cyano.

19. (Previously presented) A pharmaceutical composition comprising a cobalamin derivative according to claim 1.

20. (Previously Presented) A method of diagnosis of a neoplastic disease in a mammal comprising
(a) exposing the mammal suspected of being inflicted by a neoplastic disease or an infection to a period of a vitamin B12 – free diet, and
(b) subsequently applying a cobalamin derivative according to claim 1 carrying a diagnostic agent.

21. (Currently Amended) A method of treatment of a mammal suffering from a ~~neoplastic disease~~ melanoma comprising
(a) exposing the mammal in need of treatment to a period of a vitamin B12 – free diet, and
(b) subsequently applying a cobalamin derivative according to claim 1 carrying a therapeutic agent.

22-25. (Cancelled)

26. (Previously presented) The method of claim 20, wherein the cobalamin is effective in cancer imaging.

27. (Cancelled)